

ORIGINAL REPORT

Umbrella review of nursing interventions NIC for the treatment and prevention of suicidal behavior

Teresa Sufrate-Sorzano RN, MSc.^{1,2}  | Jesús Pérez PhD³  |
Raúl Juárez-Vela RN, MSc, PhD^{1,2}  | MaríaElena Garrote-Cámara RN, MSc.¹  |
Regina Ruiz de Viñaspre RN, MSc, PhD.¹  | Fidel Molina-Luque PhD^{4,5,6}  |
Iván Santolalla-Arnedo RN, MSc, PhD^{1,2} 

¹Research Unit on Health System Sustainability (GISSOS), Biomedical Research Center of La Rioja (CIBIR) Logroño, Logroño, Spain

²Department of Nursing, GRUPAC, University of La Rioja, Logroño, La Rioja, Spain

³Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, UK

⁴Faculty of Education, Psychology and Social Work, University of Lleida, Lleida, Spain

⁵Group for the Study of Society, Health, Education and Culture (GESEC). University of Lleida, Lleida, Spain

⁶Research Institute in Social and Territorial Development (INDEST), University of Lleida, Lleida, Spain

Correspondence

María Elena Garrote-Cámara. RN, MSc, PhDc., Clinical Research Associate, Research Unit on Health System Sustainability (GISSOS), Biomedical Research Center of La Rioja (CIBIR), 26006 Logroño, Spain.
Email: maria-elena.garrote@unirioja.es

Abstract

Purpose: The aim of this umbrella review was to determine the most effective nursing interventions for the prevention and management of suicidal behavior. In order to do so, the review identifies interventions from the Nursing Interventions Classifications taxonomy with evidence to this end.

Methods: A systematic search was conducted for systematic reviews included in the PubMed, CINAHL, Cochrane, Scopus, Web of Knowledge, and Joanna Briggs Institute databases between January 1, 2011 and May 1, 2020.

Findings: The review is made up of 21 systematic reviews. In order to carefully analyze the interventions described, these were coded into categories, grouping the interventions into those based on psychological therapy, interventions related to pharmacotherapy, interventions related to the professional relationship between health professional and patient, and interventions aimed at the general public.

Conclusions: Nursing interventions for suicide risk management are prevalent in the scientific literature and are shown to be effective for patients with suicidal ideation or suicide attempts. Prevention and treatment of suicidal behavior require the provision of coping tools, behavioral interventions that directly address suicidal thoughts and behaviors, and support through therapeutic partnerships, among others.

Implications for nursing practice: This paper synthesizes the most current evidence on the most effective Nursing Interventions Classifications interventions for the treatment and prevention of suicidal behavior. It provides nursing practitioners with a comprehensive review of the therapeutic interventions with the best evidence and is useful for the development of clinical guidelines and protocols, as well as for the development of health policies and plans.

KEYWORDS

attempted suicide, nursing interventions, risk reduction behavior, suicidal ideation, suicide

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *International Journal of Nursing Knowledge* published by Wiley Periodicals LLC on behalf of NANDA International.

INTRODUCTION

Suicidal behavior (ideation, attempt, and death by suicide) is the result of a complex interaction of factors grouped under diathesis-stress models, which are also called vulnerability-stress models. These include risk factors that can trigger suicidal behavior, such as unemployment; relationship problems; drug and alcohol abuse, and social, psychological, genetic, and educational precipitating factors, along with protective factors such as self-confidence; resilience; social integration, and lethal means restriction (Wasserman, 2001; World Health Organization, 2002). Consequently, the intervention of the nursing professional should aim to promote protective factors, reduce risk factors, and mitigate precipitating factors (Anseán, 2014; World Health Organization, 2014).

The World Health Organization – WHO (2014) has warned health authorities that suicidal behavior is a major public health problem, which is why working on its prevention and management should be a priority for public health policies worldwide. Annual figures released by the Pan American Health Organization (2020) are as high as 700,000 deaths by suicide each year worldwide, as well as more than 16 million suicide attempts (since for every suicide there are around 20 attempts). Suicide attempts are considered to be the most significant individual risk factor in the general population (World Health Organization, 2018) because these acts are repeated by up to 30% of people during the first year, with 2% subsequently committing suicide successfully in the next 5 or 10 years (Cano et al., 2009).

Globally, suicide was the fourth leading cause of death in the 15–29 age group in 2019, with more than 77% of all cases occurring in low- and middle-income countries (Pan American Health Organization, 2020). The geographical distribution of the highest suicide rates includes Slovenia, South Korea, and Lithuania, with figures in the range of 27 deaths/100,000 inhabitants. In contrast, countries such as Greece, Turkey, and South Africa have a rate of less than 4 deaths/100,000 population, with hanging and jumping being the most commonly selected methods (National Institute of Statistics, 2020).

The recently published WHO (2021) strategy for suicide prevention in all countries, “Live Life,” recommends the implementation of effective evidence-based interventions, including restricting access to the most commonly used means (pesticides, firearms, or certain medications), prevention in the general population, educating the media on the appropriate dissemination of information, developing socioemotional skills in adolescents, and early detection, follow-up, assessment and treatment of the at-risk population (World Health Organization, 2021).

Due to the multiple factors involved in the development of suicidal behavior, prevention requires the coordination and cooperation of different sectors of society, such as education, health, law enforcement, politics, and the media. Studies aimed at the prevention and treatment of suicidal behavior highlight the effectiveness of multilevel intervention in adults with significant results in reducing ideation and repetition of the behavior (Melia et al., 2020; Büscher et al., 2020; Briggs et al., 2019; Gøtzsche & Gøtzsche, 2017; Larsen et al., 2016). Cognitive behavioral therapy, individual and group support, brief contact, counseling, improved access to health services, support groups led

by health professionals, education and information, mental health promotion, community awareness, coping support, stress management, and symptom detection training have all been proposed as effective interventions (Hofstra et al., 2020; Lai et al., 2014).

In relation to the most recent publications on the effect of interventions for the prevention and management of suicidal behavior, Hofstra et al. (2020), in their systematic review which includes a meta-analysis, highlight the effectiveness of interventions such as health education and training, telephone counseling and follow-up, promotion of gatekeeper figures, and cognitive behavioral therapy. In terms of brief and contact intervention as a skilled strategy for suicide prevention, the WHO indicates that suicide rates decrease with follow-up interventions, training, education and counseling, the administration of lithium in specific pathologies, group therapy, and cognitive behavioral therapy, with the latter emerging as one of the most effective therapies in routine treatment (Gøtzsche & Gøtzsche, 2017; Lai et al., 2014; Riblet et al., 2017). Several studies report a decrease in self-harming events, following brief contact interventions including telephone or postal mail follow-up (Milner et al., 2015). New technologies employed for the promotion of contact interventions offer promising and significant results in prevention, as well as a positive impact on the promotion of coping strategies in reducing depression, distress, and self-harm (Larsen et al., 2016; Melia et al., 2020).

Professional health intervention for the care of people at risk and/or with suicidal behavior is carried out by community care teams, hospitalization and emergency services and particularly by the different units that make up the mental health care networks of the health services. From these units, the nursing professional, as part of the multidisciplinary team, develops general and specific interventions on population, patient and/or user, health education, contact, follow-up and case management, cognitive behavioral therapy, problem-solving therapy, survivor interventions, interpersonal therapy, medication administration, and crisis intervention techniques, among others (Registered Nurse’s Association of Ontario, 2009). These types of interventions carried out by nursing professionals are generally well received by the patient due to the availability, accessibility, and therapeutic relationship established between the patient and the professional (Silva et al., 2018). Nursing professionals contribute their knowledge and skills at the levels of care, management, education, and research (Galvis, 2015).

Nursing activity in relation to the intervention and prevention of suicidal behavior is fundamentally encapsulated by the ninth need described by Virginia Henderson in the Basic Needs nursing model, “Avoidance of environmental hazards,” which includes the assessment of the patient’s knowledge of environmental hazards and their corresponding prevention and intervention (Gallego et al., 2015; Henderson, 1961). The NANDA-International (NANDA-I) nursing diagnosis on the risk of suicide (00150) is defined as the situation where there is a danger of self-inflicted life-threatening injury (Ackley et al., 2021).

This study will use the nursing diagnosis Risk for suicide (00150) as a reference to determine, according to current scientific evidence, which interventions are effective in nursing practice for the prevention and treatment of suicidal behavior, identifying these in the Nursing Interventions Classification (NIC) taxonomy (Butcher et al., 2018).

Systematic reviews and meta-analyses describing interventions for the prevention and treatment of suicidal ideation and/or attempted suicidal behavior employ heterogeneous methodologies, which impede a joint perspective that would allow for the identification of the most effective interventions that nurses can implement. An umbrella review is a systematic review that synthesizes the highest level of evidence and contributes to a global and complete vision to respond to the objectives pursued, providing an assessment of the quality of the existing evidence (Aromataris et al., 2015; Biondi-Zoccai, 2016). One of the main utilities of this methodology is that a more holistic perspective on the available scientific evidence can be obtained from several systematic reviews answering the same question and with a more systematic and rigorous focus than an integrative review (Fusar-Poli & Radua, 2018). The creation of an umbrella review involves compiling the studies with the best evidence, in this case for the search for effective nursing interventions for the treatment and prevention of suicidal behavior, in order to generate a high-quality overview of the issue in question (Chambergo-Michilot et al., 2021).

OBJECTIVE

The aim of this umbrella review was to identify and gather scientific evidence on nursing interventions from the NIC taxonomy that have been shown to be effective in the prevention and treatment of suicidal behavior (Butcher et al., 2018).

The overarching review question for the study was as follows: What nursing interventions have been shown to be effective in the prevention and treatment of suicidal behavior in adults?

METHODS AND ANALYSIS

This systematic review was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. The umbrella review was registered in the International Prospective Register of Systematic Reviews (PROSPERO) with registration code CRD42020221516.

Eligibility criteria

Systematic reviews with or without meta-analyses on the effectiveness of nursing interventions on suicidal behavior in men and women over 18 years of age, in Spanish and English, and less than 10 years old. To compile evidence on interventions that could be extrapolated to the general population, research was excluded if the sample consisted of specific population groups with particular idiosyncrasies, such as military authorities, armed forces personnel, war veterans, or the prison population.

Sources of information

A systematic search was conducted of papers published between January 1, 2011 and May 1, 2020 listed in the PubMed, Cumulative Index

to Nursing and Allied Health Literature (CINAHL), Cochrane Database of Systematic Reviews (CDSR), Scopus, Institute for Scientific Information Web of Knowledge (ISI WoK), and the Joanna Briggs Institute (JBI) databases. In 2011, the European Community called on member states to include strategies focusing on the treatment and prevention of suicidal behavior in their health policies (Scheftlein, 2011). In the same year, the WHO published a protocol guide for the intervention and assessment of patients with suicidal behavior and mental disorders, among others (World Health Organization, 2011). Subsequently, at the World Health Assembly in 2013, an agreement was made to develop the "WHO Comprehensive Mental Health Action Plan 2013–2020," one of the priority goals of which was to reduce the suicide rate by 10% by 2020 (World Health Organization, 2013).

The keywords used were suicide attempted, suicide, suicidal ideation, primary prevention, secondary prevention, tertiary prevention, risks, risk assessment, risk factors, risk reduction behavior, and systematic review. These were combined with the Boolean operators and adapted specifically to each database.

Search strategy

The search strategy is shown in the file "Supplementary 1."

Selection process for studies

The inclusion of papers was assessed by two independent reviewers and they assessed the methodological quality of the included systematic reviews and meta-analyses and data extraction. Selection conflicts between reviewers were resolved by an independent third reviewer.

The process began with the initial search and exclusion of duplicate studies; the review of the title and abstract was conducted by two independent reviewers. Papers that were only selected by one reviewer were retrieved for full-text reanalysis by both reviewers. Conflict resolution was carried out by a third-party reviewer.

The search comprised a total of 3711 papers, obtained 575 from PubMed, 341 from CINAHL, 22 from CDSR, 1156 from Scopus, 1563 from WOS, and 54 from JBI. Duplicates were eliminated and 2302 results were included for analysis by title and abstract. After applying the defined inclusion and exclusion criteria, a total of 314 papers were critically analyzed. Following this analysis, 23 studies moved on to the next phase.

Data extraction procedure

The Covidence program was used as a tool for reviewing and selecting papers for review (Covidence, 2020). A form for data extraction was developed using an adapted version of the "JBI data-extraction tool." The data extracted through the extraction form and based on the research question were analyzed in order to identify and summarize the nursing interventions represented in the selected studies.




FIGURE 1 Adapted PRISMA flowchart

Risk of bias assessment (quality of reviews)

As at all stages of the process, the assessment of the quality of the papers was carried out independently by two reviewers (TSS and ISA) and in case of disagreement, a third reviewer arbitrated the decision (FML). The tool used was the “JBI Critical Appraisal Checklist for Systematic Reviews and Research Syntheses” (Joanna Briggs Institute, 2020). This instrument is based on 11 criteria that give each paper a quality rating ranging from “low” if the score is below 40%, “medium” if it scores 40–70%, and “high quality” if it scores above 70%. After two papers obtained a quality assessment of less than 40% (low quality), the research group decided to exclude them from the research, so a total of 21 reviews were finally selected (Figure 1). Regarding the results of the quality analysis, 14 documents were rated with a high-quality level, and seven with a medium-quality level. Lack of clarity in the exposition of the review question and the absence of methods to minimize errors in data extraction were the most frequent reports in the evaluation. The “Supplementary 2” document contains the complete quality analysis of the reviews included.

RESULTS

The results are presented as a narrative synthesis highlighting the main findings. The evaluated interventions were coded into categories: interventions based on psychological therapy, interventions related to pharmacotherapy, interventions related to the therapeutic relationship between health professionals and patients, and interventions aimed at the general population.

The umbrella review consists of 21 systematic reviews that were published between 2011 and 2020. The characteristics of the reviews included for analysis are shown arranged chronologically in Table 1.

Seven of the reviews included meta-analyses (Büscher et al., 2020; Hofstra et al., 2020; Briggs et al., 2019; Gøtzsche & Gøtzsche, 2017; Meerwijk et al., 2016; Milner et al., 2015; Winter et al., 2013), one paper focused on the working environment (Milner et al., 2014), and one was based on a population aged 60+ (Okolie et al., 2017). Several reviews examined the prevention of suicidal behavior through the use and employment of online tools, mobile apps, and social networks

TABLE 1 Details of the included systematic reviews

Authors	General objective	Studies	Categories of intervention that they include
Lapierre et al. (2011)	To examine the outcomes of interventions and identify successful strategies	19	Therapy based; pharmacological intervention; health professional-patient relationship
Winter et al. (2013)	To assess the current state of prevention and its effectiveness	112	Therapy based; health professional-patient relationship
Christensen et al. (2014)	Examining online suicide detection, the effectiveness of interventions, and proactive interventions after identifying people at risk from their posts	9	Therapy based; health professional-patient relationship; community
Lai et al. (2014)	Review and evaluate suicide prevention strategies available on the web	15	Therapy based; health professional-patient relationship; community
Milner et al. (2014)	Evaluate suicide prevention activities in the workplace	13	Health professional-patient relationship; community
Macedo et al. (2014)	To evaluate the effectiveness of interventions to promote resilience in adults. To assess their efficacy on suicidal behavior	13	Therapy based; health professional-patient relationship; community
Milner et al. (2015)	To synthesize the evidence regarding the effectiveness of brief contact interventions in reducing self-harm, suicide attempts, and completed suicide	12	Therapy based; health professional-patient relationship
Robinson et al. (2016)	To identify current evidence related to social media as a tool for suicide prevention	30	Therapy based; health professional-patient relationship
Meerwijk et al. (2016)	To assess whether psychosocial and behavioral interventions that address suicidal thoughts and behaviors are more effective in reducing suicides and suicide attempts than interventions that only address associated symptoms	53	Therapy based; pharmacological intervention; health professional-patient relationship
Larsen et al. (2016)	To examine the compatibility of the features of available apps with current scientific evidence on effective suicide prevention strategies	49	Therapy based; health professional-patient relationship; community
Okolie et al. (2017)	To identify and evaluate the evidence on the affectivity of interventions for older adults on preventing suicidal behavior and reducing suicidal ideation	21	Therapy based; pharmacological intervention; health professional-patient relationship; community
Gøtzsche & Gøtzsche (2017)	To assess the effectiveness of cognitive behavioral therapy in suicide prevention	20	Therapy based
Riblet et al. (2017)	To identify interventions for suicide prevention	78	Therapy-based; pharmacological intervention; health professional-patient relationship
Tighe et al. (2018)	To assess the effectiveness of acceptance and commitment therapy in reducing suicidal ideation and self-harm	5	Therapy based
Mendez-Bustos et al. (2019)	To review available scientific evidence on the effectiveness of psychotherapeutic tools designed to treat patients at risk of suicide	40	Therapy based; health professional-patient relationship; community-
Briggs et al. (2019)	To systematically review randomised controlled trials of psychoanalytic and psychodynamic psychotherapies for suicide attempts and self-harm	12	Therapy based; health professional-patient relationship
Dt'Anci et al. (2019)	To assess risks/benefits of suicide prevention interventions	23	Therapy based; pharmacological intervention.
Hoffberg et al. (2019)	To review the clinical effectiveness of psychotherapeutic tools designed to treat patients at risk of suicide	40	Therapy based; health professional-patient relationship
Melia et al. (2020)	To examine the effectiveness of available mobile technology tools for the prevention of any suicidal tendencies	7	Health professional-patient relationship
Büscher et al. (2020)	To study whether internet-based self-help interventions are associated with a reduction in suicidal ideation	6	Therapy based
Hofstra et al. (2020)	Evaluating the effect of suicide prevention interventions	16	Therapy based; health professional-patient relationship; community

(Büscher et al., 2020; Melia et al., 2020; Larsen et al., 2016; Robinson et al., 2016; Christensen et al., 2014; Lai et al., 2014).

Six reviews examine which interventions are effective for the prevention of suicidal behavior (Hoffberg et al., 2019; Hofstra et al., 2020; D'Anci et al., 2019; Méndez-Bustos et al., 2019; Riblet et al., 2017; Lapierre et al., 2011) and more specifically, some work focused on evaluating the functionality of specific interventions in the prevention of suicidal behaviors such as the development of resilience in the adult population (Macedo et al., 2014), brief contact (Milner et al., 2015), or acceptance and commitment therapy (Tighe et al., 2018).

Interventions based on psychological therapies

Under the category of interventions based on psychological therapies, dialectical behavioral therapy, cognitive behavioral therapy, problem solving and adjustment therapy, psychodynamic therapy, psychoanalytic therapy, acceptance and commitment therapy, and family, interpersonal, or group therapy are observed as effective interventions (Hofstra et al., 2020; Briggs et al., 2019; D'Anci et al., 2019; Hoffberg et al., 2019; Méndez-Bustos et al., 2019; Tighe et al., 2018; Gøtzsche & Gøtzsche, 2017; Okolie et al., 2017; Riblet et al., 2017; Larsen et al., 2016; Meerwijk et al., 2016; Milner et al., 2015; Winter et al., 2013; Lapierre et al., 2011).

Interventions focusing on pharmacological treatment

In the review, the health interventions focusing on pharmacological treatment that stand out for their effectiveness are education and training on the prescribed treatment; long-term control and monitoring of pharmacological guidelines; information and education on the most frequent side effects; adequate monitoring of the patient with prescriptions for ketamine and lithium; promotion of adherence to prescribed guidelines and pharmacological treatment for affective disorders (Okolie et al., 2017; Riblet et al., 2017; Meerwijk et al., 2016; Lapierre et al., 2011).

Interventions related to the relationship between the health professional and the patient

Health interventions, related to the broader health professional-patient relationship, would include active patient follow-up and the establishment of follow-up meetings/consultations, brief intervention (via telephone, email, or mail), early detection and treatment of depression and its symptoms, training, and counseling in self-care, acceptance, social and coping skills training, mood monitoring and promotion of resilience (Hofstra et al., 2020; Melia et al., 2020; Briggs et al., 2019; Hoffberg et al., 2019; Méndez-Bustos et al., 2019; Okolie et al., 2017; Riblet et al., 2017). Stress management, strengthening protective factors, crisis contact and support, mood and emotion management, therapeutic alliance, support group membership, and the development

of a personalized care plan are also reported as effective interventions (Larsen et al., 2016; Meerwijk et al., 2016; Robinson et al., 2016; Milner et al., 2015; Christensen et al., 2014; Lai et al., 2014; Macedo et al., 2014; Milner et al., 2014; Winter et al., 2013; Lapierre et al., 2011). For the implementation of these interventions, scheduled nursing consultations, telephone, and new technology-based follow-ups and consultations, and supportive home visits through assertive community therapies have been shown to be effective (Büscher et al., 2020; Robinson et al., 2016; Christensen et al., 2014; Lai et al., 2014; Macedo et al., 2014). Because of the strong relationship between suicide and hopelessness (Ribeiro et al., 2018), interventions for more acute contexts may include providing hope, managing self-harming behavior, limit setting, and crisis intervention or even first aid (Butcher et al., 2018).

Interventions targeting the general public

Interventions related to the general population (society/community) for the treatment and prevention of suicidal behavior include raising awareness and understanding of the problem, facilitating access to health services, reducing access to the most commonly used means, promoting the role of community gatekeepers, promoting mental health and reducing isolation. In addition, they encompass the production and distribution of standardized, nonsensationalized, truthful information through social networks, media, and websites, as well as the early identification of harmful content (Hofstra et al., 2020; Méndez-Bustos et al., 2019; Okolie et al., 2017; Christensen et al., 2014; Larsen et al., 2016; Lai et al., 2014; Macedo et al., 2014; Milner et al., 2014). In this context, the interventions of the NIC taxonomy that are related to the community aspects are as follows: environmental management – safety; health education in general and particularly in the prevention of substance abuse; establishment of consultations or visits that promote mental health; listening visits; exchange of health care information; improvement of access to health information; and training in first aid. In addition, and specifically in the community closest to the patient, increasing family and social support systems, and enhancing socialization are effective interventions (Butcher et al., 2018).

Table 2 shows specifically coded interventions from the NIC taxonomy that are most effective for the correct management of patients with suicidal behavior.

DISCUSSION

This review has identified and compiled the scientific evidence that addresses the prevention, management, and treatment of suicidal behavior using nursing interventions from the NIC taxonomy so that the objective set by the research team has been achieved.

Due to the magnitude of the serious public health problem of suicide, the reduction of suicidal behavior must become a priority in social, political, media and health agendas, where greater

TABLE 2 Interventions codified in the NIC taxonomy which are most effective for the proper management of patients exhibiting suicidal behavior

- 1460. Progressive muscle relaxation:** Facilitating the tensing and releasing of successive muscle groups while attending to the resulting differences in sensation
- 2300. Medication administration:** Preparing, giving, and evaluating the effectiveness of prescription and nonprescription drugs
- 2395. Medication reconciliation:** Comparison of the patient's home medications with the admission, transfer, and/or discharge orders to ensure accuracy and patient safety
- 2380. Medication management:** Facilitation of safe and effective use of prescription and over-the-counter drugs
- 4340. Assertiveness training:** Assistance with the effective expression of feelings, needs, and ideas while respecting the rights of others
- 4350. Behavior management:** Helping a patient to manage negative behavior
- 4352. Behavior modification:** Provision of a therapeutic milieu that safely accommodates the patient's attention deficit and/or overactivity while promoting optimal function
- 4354. Behavior management: Self-harm:** Assisting the patient to decrease or eliminate self-mutilating or self-abusive behaviors
- 4360. Behavior modification:** Promoting behavior change.
- 4380. Setting limits:** Establishing the parameters of desirable and acceptable patient behavior.
- 4410. Establishment of common objectives:** Collaboration with the patient to identify and prioritize care goals and then develop a plan to achieve these goals.
- 4420. Patient contracting:** Negotiating an agreement with an individual that reinforces a specific behavior change
- 4480. Self-responsibility facilitation:** Encouraging a patient to assume more responsibility for own behavior
- 4500. Substance use prevention:** Prevention of an alcoholic or drug use lifestyle
- 4700. Cognitive restructuring:** Challenging a patient to alter distorted thought patterns and view self and the world more realistically
- 4860. Reminiscence therapy:** Using the recall of past events, feelings, and thoughts to facilitate pleasure, quality of life, or adaptation to present circumstances
- 4920. Active listening:** Attending closely to and attaching significance to a patient's verbal and nonverbal messages
- 5000. Complex relationship building:** Establishing a therapeutic relationship with a patient to promote insight and behavioral change
- 5020. Conflict mediation:** Facilitation of constructive dialogue between opposing parties with the aim of resolving conflicts in a mutually acceptable manner.
- 5100. Socialization enhancement:** Facilitation of another person's ability to interact with others
- 5230. Coping enhancement:** Facilitation of cognitive and behavioral efforts to manage perceived stressors, changes, or threats that interfere with meeting life demands and roles
- 5240. Counseling:** Use of an interactive helping process focusing on the needs, problems, or feelings of the patient and significant others to enhance or support coping, problem solving, and interpersonal relationships
- 5250. Decision-making support:** Providing information and support for a patient who is making a decision regarding health care
- 5270. Emotional support:** Provision of reassurance, acceptance, and encouragement during times of stress
- 5300. Guilt work facilitation:** Helping another to cope with painful feelings of actual or perceived responsibility
- 5310. Hope inspiration:** Enhancing the belief in one's capacity to initiate and sustain actions
- 5326. Life skills enhancement:** Developing an individual's ability to independently and effectively deal with the demands and challenges of everyday life
- 5328. Listening visits:** Empathic listening to genuinely understand an individual's situation and work collaboratively over a number of home visits to identify and generate solutions to reduce depressive symptoms
- 5330. Mood management:** Providing for safety, stabilization, recovery, and maintenance of a patient who is experiencing a dysfunctionally depressed or elevated mood
- 5380. Security enhancement:** Intensifying a patient's sense of physical and psychological safety
- 5390. Self-awareness enhancement:** Assisting a patient to explore and understand his/her thoughts, feelings, motivations, and behaviors
- 5395. Self-efficacy enhancement:** Strengthening an individual's confidence in his/her ability to perform a healthy behavior
- 5400. Self-esteem enhancement:** Assisting a patient to increase his or her personal judgment of self-worth
- 5430. Support group:** Use of a group setting to provide emotional support and health-related information to its members.
- 5440. Support system enhancement:** Facilitation of support to patient by family, friends, and community
- 5450. Therapy group:** Application of psychotherapeutic techniques to a group, including the utilization of interactions between members of the group

(Continues)

TABLE 2 (Continued)

5460. Touch: Providing comfort and communication through purposeful tactile contact
5480. Values clarification: Assisting another to clarify her/his own values in order to facilitate effective decision-making
5510. Health education: Developing and providing instruction and learning experiences to facilitate voluntary adaptation of behavior conducive to health in individuals, families, groups, or communities
5515. Health literacy enhancement: Assisting individuals with limited ability to obtain, process, and understand information related to health and illness
5616. Teaching: prescribed medication: Preparing a patient to safely take prescribed medications and monitor for their effects
5880. Calming technique: Reducing anxiety in patient experiencing acute distress
5960. Meditation facilitation: Facilitating a person to alter his/her level of awareness by focusing specifically on an image or thought
6040. Relaxation therapy: Use of techniques to encourage and elicit relaxation for the purpose of decreasing undesirable signs and symptoms such as pain, muscle tension, or anxiety
6160. Crisis intervention: Use of short-term counseling to help the patient cope with a crisis and resume a state of functioning comparable to or better than the precrisis state
6240. First aid: Providing immediate care for minor burns, injuries, poisoning, bites, and stings
6340. Suicide prevention: Reducing the risk for self-inflicted harm with intent to end life
6486. Environmental management – safety: Monitoring and manipulation of the physical environment to promote safety
6520. Health screening: Detecting health risks or problems by means of history, examination, and other procedures
6610. Risk identification: Analysis of potential risk factors, determination of health risks, and prioritization of risk reduction strategies for an individual or group
7100. Family integrity promotion: Promotion of family cohesion and unity
7110. Family involvement promotion: Facilitating participation of family members in the emotional and physical care of the patient
7130. Family process maintenance: Minimization of family process disruption effects
7140. Family support: Promotion of family values, interests, and goals
7150. Family therapy: Assisting family members to move their family towards a more productive way of living
7200. Normalization promotion: Assisting parents and other family members of children with chronic illnesses or disabilities in providing normal life experiences for their children and families
7800. Quality monitoring: Systematic collection and analysis of an organization's quality indicators for the purpose of improving patient care
7910. Consultation: Using expert knowledge to work with those who seek help in problem solving to enable individuals, families, groups, or agencies to achieve identified goals
7960. Health care information exchange: Providing patient care information to other health professionals
8020. Multidisciplinary care conference: Planning and evaluating patient care with health professionals from other disciplines
8100. Referral: Arrangement for services by another care provider or agency
8180. Telephone consultation: Eliciting patient's concerns, listening, and providing support, information, or teaching in response to patient's stated concerns, over the telephone
8190. Telephone follow-up: Providing results of testing or evaluating patient's response and determining potential for problems as a result of previous treatment, examination, or testing, over the telephone
8340. Resiliency promotion: Assisting individuals, families, and communities in development, use, and strengthening of protective factor to be used in coping with environmental and societal stressors

awareness and training of professionals are needed, through knowledge of evidence-based interventions. This article identifies the most effective interventions from the NIC taxonomy described for the prevention and treatment of patients presenting with suicidal ideation and behavior. In order to get a broad and comprehensive view of existing interventions supported by the best evidence, an extensive search was conducted, including systematic reviews and meta-analyses. An assessment of the methodological quality of the included systematic reviews and meta-analyses by two independent reviewers determined a high-quality rating for most of the studies. Taking into account the concepts

that integrate the holistic nursing metaparadigm (person, health, environment, care), as well as the different integral models of nursing care such as the critical-holistic model of Miotto Wright (Wright, 2000), intervention against suicidal ideation and the prevention of suicidal behavior offers the nursing discipline a wide field of intervention. The results of this study confirm the efficacy of holistic (bio-psycho-social) interventions, based on psychological therapy, pharmacotherapy, interventions related to the relationship between the health professional and the patient, and interventions aimed at the general population.

NIC interventions based on psychological therapies for the management and reduction of suicide risk are prevalent in the scientific literature and are shown to be effective for patients with suicidal ideation or previous suicide attempts (D'Anci et al., 2019; Gøtzsche & Gøtzsche, 2017; Lai et al., 2014; Meerwijk et al., 2016; Méndez-Bustos et al., 2019; Milner et al., 2014). In this line, P. C. Gøtzsche and P. K. Gøtzsche associated interventions based on modifications of the patient's thought processes, improvement of self-confidence, and cognitive restructuring with a 50% reduction in the repetition of suicidal behavior. Interventions to build resilience, and improve self-awareness, self-esteem and coping have been shown to be effective in reducing suicidal risk (Pérez et al., 2021; Revuelta et al., 2016). After these interventions, patients' expression improved self-acceptance and were able to recognize some positive personal characteristics; several studies report a significant improvement in the Rosenberg Self-Esteem Scale after these NIC interventions (Rodríguez et al., 2014). The use of self-help interventions, aimed at working on self-esteem and coping strategies, based on new information technologies (chats, social networks, apps, email, and so on) are associated with significant reductions in suicidal ideation when developed on the basis of scientific evidence and best practices (Büscher et al., 2020; Christensen et al., 2014; Lai et al., 2014; Larsen et al., 2016; Melia et al., 2020; Robinson et al., 2016). In this area, some studies have shown the development of these techniques in online formats using new technologies and tools to be effective, including mobile applications, group chats, and social networks (Büscher et al., 2020; Robinson et al., 2016; Christensen et al., 2014; Lai et al., 2014; Macedo et al., 2014).

Patients with suicidal ideation and/or behavior often require pharmacological treatment, especially when suicidal ideation and/or behavior is related to the presence of a mental disorder (Barbero et al., 2010; Sansaloni et al., 2020; Rodríguez et al., 2014). NIC nursing interventions related to medication management, including patient education (teaching of prescribed medication) on the use of psychotropic drugs, appropriate follow-up by nursing professionals, adherence, and side effect management showed significant evidence for the prevention and treatment of suicidal behavior (D'Anci et al., 2019; Lapierre et al., 2011; Meerwijk et al., 2016; Okolie et al., 2017; Riblet et al., 2017). Mood regulators, such as lithium, are among the most commonly used drugs in these preventive programs (D'Anci et al., 2019).

Interventions related to the nurse/patient relationship should be singled out as high-quality preventive interventions by assessing the active follow-up of patients and enabling the nursing practice to develop NIC interventions such as health counseling and education, decision support, values clarification, emotional support, mood management, active listening, and contact. Several studies link brief contact interventions to a reduction in suicide attempts (Milner et al., 2015; Riblet et al., 2017). The effectiveness of these interventions has been demonstrated both for direct contact in consultation or at home, and for contact by telephone, email, or post (Büscher et al., 2020; Christensen et al., 2014; Melia et al., 2020; Riblet et al., 2017; Robinson et al., 2016; Consejería de Sanidad y Dependencia, 2008). The scientific literature has linked contact interventions where emotional support is provided to the reduction of self-harm related to suicidal

ideation (Briggs et al., 2019; Hoffberg et al., 2019; Méndez-Bustos et al., 2019; Winter et al., 2013). The therapeutic nurse/patient relationship is a protective element against suicidal behavior (Briggs et al., 2019; Lapierre et al., 2011; Riblet et al., 2017); from this supportive relationship, it is possible to help maintain positive family processes and promote family integration. Some authors have emphasized the preventive role of the nuclear family. The family context can become a positive source in terms of self-control, verbal conflict resolution, counseling, emotional support, monitoring, and protection from self-destructive and risky behaviors (Becoña Iglesias, 2001; Sánchez et Carrillo, 2002; Barbero et al., 2010; Villa et Hermida, 2001).

In terms of community intervention, in the NIC taxonomy, we can highlight the identification and management of risks in the environment (environmental management). In this regard, restricting access to the most widely used lethal methods is one of the most prevalent preventive measures; however, efforts should be directed towards exploring new methods which could replace them, with the aim of working proactively (Méndez-Bustos et al., 2019; Okolie et al., 2017; Christensen et al., 2014; Larsen et al., 2016; Lai et al., 2014). Health education and improved access to health information are key preventive elements. It is a priority to "break the silence," the traditional obscurantism, and to break with false myths, thinking and beliefs in relation to suicidal behavior, proposing actions to raise awareness and promote mental health from nursing consultations and advising the media in the dissemination of campaigns and news (Brito et al., 2020; Revuelta et al., 2016; Galán et al., 2010). New communication technologies, including social networks, chat rooms, online platforms, and mobile applications, are now key tools for community awareness raising. The reach, anonymity, instantaneity, and accessibility of these new information and communication technologies allow them to act as key elements alongside traditional media (Larsen et al., 2016; Macedo et al., 2014; Robinson et al., 2016).

Based on the latest edition of the NIC catalog of interventions (Butcher et al., 2018), this review identifies the various interventions that have been shown to be effective for individual and group work with the patient, family, and community. Prevention and treatment of suicidal behavior require comprehensive management in which patients, family, and community are provided with coping tools; behavioral interventions that directly address suicidal thoughts and behaviors; emotional management through psychological therapies; conventional therapies together with new information and communication technologies; support through therapeutic alliances and brief contact interventions; support for adherence to treatment, environmental management, and health education for patients and community.

LIMITATIONS

The research team worked within a time range of 10 years and used the latest published version of the NIC taxonomy, but the review focuses only on the most effective and specific NIC interventions for the prevention and treatment of suicidal behaviors, so there may be a bias of

interventions available to the nurse that is not described in this paper, as there is also learning from negative results.

CONCLUSION

This paper offers a highly methodological synthesis that identifies the most effective NIC interventions for the treatment and prevention of suicidal behavior. It provides a thorough and comprehensive review of the best evidence-based therapeutic interventions and can serve as a basis for clinical intervention guidelines and protocols and specific suicide prevention plans. These results enable nursing professionals to use the evidence to make decisions in their daily practice.

IMPLICATIONS FOR NURSING PRACTICE, RESEARCH, AND EDUCATION

This umbrella review synthesizes the most up-to-date evidence on the most effective NIC interventions for the treatment and prevention of suicidal behavior. It provides nursing practice with a review of the therapeutic interventions with the best evidence, being useful for the development of guidelines and protocols in clinical and care settings, as well as providing the up-to-date and evidence-based documentation that is necessary for the development of health policies and plans focused on the prevention and management of suicidal behaviors.

The standardized nursing interventions that have been detailed in this work allow the care provided to the patient, family, and community to be recorded and evaluated in a universal way, and thus to assess the impact on the needs and health of the units to be treated. Nursing professionals working in the field of research use standardized nursing language that facilitates the study and analysis of the quality and efficacy of the care provided. Therefore, the results obtained in this review can be used as quality indicators and can easily be compared with interventions and care provided worldwide. In the same way, these results synthesize the most current and effective scientific evidence, making it a necessary tool in the educational field for training future nursing professionals.

ACKNOWLEDGMENTS

The research team would like to thank the Research Group on Care (GRUPAC) of the University of La Rioja and the Research Group on Sustainability of the Health System (GISOSS) of the Biomedical Research Centre of La Rioja (CIBIR) for their support and collaboration. There was no financial support for this article.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

Conceptualization: Teresa Sufrate-Sorzano and María Elena Garrote-Cámara.

Methodology: Teresa Sufrate-Sorzano, Jesús Pérez, María Elena Garrote-Cámara, and Regina Ruiz de Viñaspre.

Software: Raúl Juárez-Vela and Fidel Molina-Luque.

Validation: Jesús Pérez, María Elena Garrote-Cámara, Iván Santolalla-Arnedo and Raúl Juárez-Vela.

Formal análisis: Teresa Sufrate Sorzano, Iván Santolalla-Arnedo, and Fidel Molina-Luque.

Investigation: Teresa Sufrate-Sorzano and María Elena Garrote-Cámara.

Resources: Jesús Pérez, Iván Santolalla-Arnedo, Raúl Juárez-Vela, Fidel Molina-Luque, and Regina Ruiz de Viñaspre.

Data curation: Iván Santolalla-Arnedo, Raúl Juárez-Vela, Fidel Molina-Luque, and Regina Ruiz de Viñaspre.

Writing—original draft preparation: Teresa Sufrate-Sorzano, Jesús Pérez, and María Elena Garrote-Cámara.

Writing—review and editing: Iván Santolalla-Arnedo, Raúl Juárez-Vela, Fidel Molina-Luque, and Regina Ruiz de Viñaspre.

Visualization: Teresa Sufrate-Sorzano and María Elena Garrote-Cámara.

Supervision: Jesús Pérez, Iván Santolalla-Arnedo, Raúl Juárez-Vela, Fidel Molina-Luque, and Regina Ruiz de Viñaspre.

Project administration: Teresa Sufrate-Sorzano.

All authors have read and agreed to the published version of the manuscript.

ETHICS STATEMENT

Formal ethical approval is not required for this umbrella review under Spanish law, as no primary data are collected, and no patient intervention is involved.

PROSPERO Registration number: CRD42020221516.

ORCID

Teresa Sufrate-Sorzano RN, MSc.  <https://orcid.org/0000-0003-3756-9914>

Jesús Pérez PhD  <https://orcid.org/0000-0003-0740-190X>

Raúl Juárez-Vela RN, MSc, PhD  <https://orcid.org/0000-0003-3597-2048>

María Elena Garrote-Cámara RN, MSc.  <https://orcid.org/0000-0002-7504-4070>

Regina Ruiz de Viñaspre RN, MSc, PhD.  <https://orcid.org/0000-0003-0345-8028>

Fidel Molina-Luque PhD  <https://orcid.org/0000-0001-5278-2794>

Iván Santolalla-Arnedo RN, MSc, PhD  <https://orcid.org/0000-0001-6705-7122>

REFERENCES

- Anseán, A. (2014). *Spanish Foundation for Mental Health for the Prevention of Mental Disorders and Suicide, Suicides*. La Rioja Government.
- Aromataris, E., Fernandez, R., Godfrey, C. M., Holly, C., Khalil, H., & Tungpunkom, P. (2015). Summarizing systematic reviews: Methodological development, conduct and reporting of an umbrella review approach. *International Journal of Evidence-Based Healthcare*, 13(3), 132–140. <https://doi.org/10.1097/XEB.0000000000000055>

- Barbero, J. J., Rocamora, C. R., García, I. D., García, M. P., & Graupera, J. X. (2010). Intervenciones enfermeras en trastornos duales: Adicciones relacionadas con trastornos de conducta en la infancia. *Enfermería Global*, 9(2), 1–12.
- Becoña Iglesias, E. (2001). Factores de riesgo y protección familiar para el uso de drogas. In Plan Nacional sobre Drogas (Ed.) *Intervención familiar en la prevención de las drogodependencias*. 113–140.
- Biondi-Zoccai, G. (2016). *Umbrella reviews. Evidence synthesis with overviews of reviews and meta-epidemiologic studies*. Springer International.
- Briggs, S., Netuveli, G., Gould, N., Gkaravella, A., Gluckman, N. S., Kangogere, P., Farr, R., Goldblatt, M. J., & Lindner, R. (2019). The effectiveness of psychoanalytic/psychodynamic psychotherapy for reducing suicide attempts and self-harm: Systematic review and meta-analysis. *The British Journal of Psychiatry: The Journal of Mental Science*, 214(6), 320–328. <https://doi.org/10.1192/bjp.2019.33>
- Brito, P. R., Bazán Herrero, A., Castrillo García, F. B., Díez Álvarez, A. B., García García, R., García Fernante, M., Gómez Feliz, N., González García, M., González Pérez, F., Gutiérrez Fernández, M. J., Iglesias Suárez, M., Iglesias Valbuena, M. M., Peláez Nava, A., Riesgo Laviana, A., Sierra Gayo, M. B., & Suárez Iglesias, M. J. (2020). Priorización de diagnósticos, criterios de resultado e intervenciones enfermeras psicosociales en el contexto de un programa formativo en lenguaje de cuidados. *Ene*, 14(3), e14302. <https://doi.org/10.4321/s1988-348x2020000300002>
- Büscher, R., Torok, M., Terhorst, Y., & Sander, L. (2020). Internet-based cognitive behavioral therapy to reduce suicidal ideation: A systematic review and meta-analysis: A systematic review and meta-analysis. *JAMA Network Open*, 3(4), e203933. <https://doi.org/10.1001/jamanetworkopen.2020.3933>
- Butcher, H. K., Bulechek, G. M., Dochterman, J. M. M., & Wagner, C. M. (2018). *Nursing interventions classification (NIC)-E-Book*. Elsevier Health Sciences.
- Cano, P., Gutierrez, C., & Nizama, M. (2009). Tendency to violence and suicidal ideation in school-aged youths in a city of the Peruvian Amazon. *Revista Peruana de Medicina Experimental y Salud Pública*, 26(2), 175–181.
- Chambergo-Michilot, D., Díaz-Barrera, M. E., & Benites-Zapata, V. A. (2021). Revisiones de alcance, revisiones paraguas y síntesis enfocada en revisión de mapas: Aspectos metodológicos y aplicaciones. *Revista peruana de medicina experimental y salud pública*, 38(1), 136–142. <https://doi.org/10.17843/rpmpesp.2021.381.6501>
- Christensen, H., Batterham, P. J., & O'Dea, B. (2014). E-health interventions for suicide prevention. *International Journal of Environmental Research and Public Health*, 11(8), 8193–8212. <https://doi.org/10.3390/ijerph110808193>
- Covidence [program]. (2020). Veritas Health Innovation. www.covidence.org
- D'Anci, K. E., Uhl, S., Giradi, G., & Martin, C. (2019). Treatments for the prevention and management of suicide: A systematic review: A systematic review. *Annals of Internal Medicine*, 171(5), 334–342. <https://doi.org/10.7326/M19-0869>
- Fusar-Poli, P., & Radua, J. (2018). Ten simple rules for conducting umbrella reviews. *Evidence-Based Mental Health*, 21(3), 95–100.
- Galán, A. M. R., González, M. D. C. P., & Ferrón, S. B. (2010). Plan de cuidados del paciente con trastorno de personalidad dependiente. *NURE investigación: Revista Científica de enfermería*, 48, 4.
- Gallego, R., Díez, J., & López, A. (2015). Metodología Enfermera: Lenguajes estandarizados. *Ene*, 9, 40.
- Galvis, M. A. (2015). Teorías y modelos de enfermería usados en la enfermería psiquiátrica. *Revista CUIDARTE*, 6(2), 1108. <https://doi.org/10.15649/cuidarte.v6i2.172>
- Gøtzsche, P. C., & Gøtzsche, P. K. (2017). Cognitive behavioural therapy halves the risk of repeated suicide attempts: Systematic review. *Journal of the Royal Society of Medicine*, 110(10), 404–410. <https://doi.org/10.1177/0141076817731904>
- Henderson, V. (1961). *Principios Básicos de los Cuidados de Enfermería*. Publicación científica no. 57. Organización Mundial de la Salud.
- Hoffberg, A. S., Stearns-Yoder, K. A., & Brenner, L. A. (2019). The effectiveness of crisis line services: A systematic review. *Frontiers in Public Health*, 7, 399. <https://doi.org/10.3389/fpubh.2019.00399>
- Hofstra, E., van Nieuwenhuizen, C., Bakker, M., Özgül, D., Elfeddali, I., de Jong, S. J., & van der Feltz-Cornelis, C. M. (2020). Effectiveness of suicide prevention interventions: A systematic review and meta-analysis. *General Hospital Psychiatry*, 63, 127–140. <https://doi.org/10.1016/j.genhosppsych.2019.04.011>
- Joanna Briggs Institute. (2020). The Joanna Briggs Institute critical appraisal tools for use in JBI systematic reviews. Checklist for systematic reviews and research syntheses. <http://joannabriggs.org/research/critical-appraisal-tools.html>
- Lai, M. H., Maniam, T., Chan, L. F., & Ravindran, A. V. (2014). Caught in the web: A review of web-based suicide prevention. *Journal of Medical Internet Research*, 16(1), e30. <https://doi.org/10.2196/jmir.2973>
- Lapierre, S., Erlangsen, A., Waern, M., De Leo, D., Oyama, H., Scocco, P., Gallo, J., Szanto, K., Conwell, Y., Draper, B., & Quinnett, P., & International Research Group for Suicide among the Elderly. (2011). A systematic review of elderly suicide prevention programs. *Crisis*, 32(2), 88–98. <https://doi.org/10.1027/0227-5910/a000076>
- Larsen, M. E., Nicholas, J., & Christensen, H. (2016). A systematic assessment of smartphone Tools for suicide prevention. *Plos ONE*, 11(4), e0152285. <https://doi.org/10.1371/journal.pone.0152285>
- Macedo, T., Wilhelm, L., Gonçalves, R., Coutinho, E. S. F., Vilete, L., Figueira, I., & Ventura, P. (2014). Building resilience for future adversity: A systematic review of interventions in non-clinical samples of adults. *BMC Psychiatry [Electronic Resource]*, 14(1), 227. <https://doi.org/10.1186/s12888-014-0227-6>
- Meerwijk, E. L., Parekh, A., Oquendo, M. A., Allen, I. E., Franck, L. S., & Lee, K. A. (2016). Direct versus indirect psychosocial and behavioural interventions to prevent suicide and suicide attempts: A systematic review and meta-analysis. *The Lancet Psychiatry*, 3(6), 544–554. [https://doi.org/10.1016/S2215-0366\(16\)00064-X](https://doi.org/10.1016/S2215-0366(16)00064-X)
- Melia, R., Francis, K., Hickey, E., Bogue, J., Duggan, J., O'Sullivan, M., & Young, K. (2020). Mobile health technology interventions for suicide prevention: Systematic review. *JMIR mHealth and uHealth*, 8(1), e12516. <https://doi.org/10.2196/12516>
- Méndez-Bustos, P., Calati, R., Rubio-Ramírez, F., Olié, E., Courtet, P., & Lopez-Castroman, J. (2019). Effectiveness of psychotherapy on suicidal risk: A systematic review of observational studies. *Frontiers in Psychology*, 10, 277. <https://doi.org/10.3389/fpsyg.2019.00277>
- Milner, A., Page, K., Spencer-Thomas, S., & Lamotagne, A. D. (2014). Workplace suicide prevention: A systematic review of published and unpublished activities. *Health Promotion International*, 30(1), 29–37. <https://doi.org/10.1093/heapro/dau08>
- Milner, A. J., Carter, G., Pirkis, J., Robinson, J., & Spittal, M. J. (2015). Letters, green cards, telephone calls and postcards: Systematic and meta-analytic review of brief contact interventions for reducing self-harm, suicide attempts and suicide. *The British Journal of Psychiatry: The Journal of Mental Science*, 206(3), 184–190. <https://doi.org/10.1192/bjp.bp.114.147819>
- National Institute of Statistics. (2020). Deaths from causes, gender and age groups. National Institute of Statistics. <http://www.ine.es/jaxiT3/Datos.htm?t=7947>
- Okolie, C., Dennis, M., Simon Thomas, E., & John, A. (2017). A systematic review of interventions to prevent suicidal behaviors and reduce suicidal ideation in older people. *International Psychogeriatrics*, 29(11), 1801–1824. <https://doi.org/10.1017/S1041610217001430>
- Pan American Health Organization. (2020). Suicide prevention. Pan American Health Organization. <https://www.paho.org/en/topics/suicide-prevention>
- Pérez, S., Gargallo, A., Aranda Hernández, M., & Sánchez, F. (2021). Plan de cuidados de enfermería en salud mental: Ideación suicida, caso clínico. *Revista Sanitaria de Investigación (RSI)*.

- <https://revistasanitariadeinvestigacion.com/plan-de-cuidados-de-enfermeria-en-salud-mental-ideacion-suicida-caso-clinico/>
- Registered Nurse's Association of Ontario. (2009). Guía de buenas prácticas en enfermería. Valoración y cuidado de los adultos en riesgo de ideación y conducta suicida. Investén-Isciii.
- Revuelta, J. F. R., Vives, E. R., & Blanco, L. (2016). Plan de cuidados en Enfermería del Trabajo Caso práctico: Trabajador con ideación suicida. *Revista Enfermería Del Trabajo*, 6(3), 96–114.
- Ribeiro, J. D., Huang, X., Fox, K. R., & Franklin, J. C. (2018). Depression and hopelessness as risk factors for suicide ideation, attempts and death: Meta-analysis of longitudinal studies. *The British Journal of Psychiatry: The Journal of Mental Science*, 212(5), 279–286. <https://doi.org/10.1192/bjp.2018.27>
- Riblet, N. B. V., Shiner, B., Young-Xu, Y., & Watts, B. V. (2017). Strategies to prevent death by suicide: Meta-analysis of randomised controlled trials. *The British Journal of Psychiatry: The Journal of Mental Science*, 210(6), 396–402. <https://doi.org/10.1192/bjp.bp.116.187799>
- Robinson, J., Cox, G., Bailey, E., Hetrick, S., Rodrigues, M., Fisher, S., & Herrman, H. (2016). Social media and suicide prevention: A systematic review: Suicide prevention and social media. *Early Intervention in Psychiatry*, 10(2), 103–121. <https://doi.org/10.1111/eip.12229>
- Rodríguez, Á. M. M., Jiménez, R. P. M., Santana, C. H., & Artiles, F. J. A. (2014). Plan de cuidados de enfermería a una paciente con esquizofrenia y baja autoestima crónica. *Revista Enfermería Docente*, 102, 35–41.
- Sansaloni, M., Sanchez, V., & Ibañez, L. (2020). Plan de cuidados en el paciente psiquiátrico en urgencias. *Revista Electrónica de portales Médicos*. <https://www.revista-portalesmedicos.com/revista-medica/plan-de-cuidados-en-el-paciente-psiquiatrico-en-urgencias/>
- Sánchez, J. P. E., & Carrillo, F. J. M. (2002). Factores familiares, comportamientos perturbadores y drogas en la adolescencia. In Plan Nacional sobre Drogas (Ed.). *Intervención familiar en la prevención de drogodependencias*. 25–55.
- Consejería de Sanidad y Dependencia. (2008). *Planes de cuidados enfermeros en salud mental. Cuaderno técnico de salud mental*. Junta de Extremadura. Servicio Extremeño de Salud.
- Schefflein, J. (2011). The European pact for mental health and well-being. *Psychiatrische Praxis*, 38(S 01), <https://doi.org/10.1055/s-0031-1277729>
- Silva, A., Dawson, H., Gawarammana, A., Tennakoon, I. B., & Rajapakse, S. (2018). Study protocol: A pilot randomized controlled trial to evaluate the acceptability and feasibility of a counseling intervention, delivered by nurses, for those who have attempted self-poisoning in Sri Lanka. *Pilot Feasibility Studies*, 4(1), 1–9.
- Tighe, J., Nicholas, J., Shand, F., & Christensen, H. (2018). Efficacy of acceptance and commitment therapy in reducing suicidal ideation and deliberate self-harm: Systematic review. *JMIR Mental Health*, 5(2), e10732. <https://doi.org/10.2196/10732>
- Villa, S., & Hermida, R. (2001). Factores de riesgo familiares para el uso de las drogas: un estudio empírico español. In Plan Nacional sobre Drogas (Ed.). *Intervención familiar en la prevención de drogodependencias*. 57–112.
- Wasserman, D. (2001). *Suicide: An unnecessary death*. Martin Dunitz.
- Winter, D., Bradshaw, S., Bunn, F., & Wellsted, D. (2013). A systematic review of the literature on counselling and psychotherapy for the prevention of suicide: 1. Quantitative outcome and process studies. *Counselling and Psychotherapy Research*, 13(3), 164–183. <https://doi.org/10.1080/14733145.2012.761717>
- World Health Organization. (2002). *Multisite intervention study on suicidal behaviours-SUPRE-MISS: Protocol of SUPRE-MISS*. <https://iris.paho.org/handle/10665.2/45518>
- World Health Organization. (2011). *Intervention guide for mental, neurological and substance use disorders in non-specialized health settings: mental health Gap Action Programme*. <https://www.who.int/publications/i/item/9789241549790>
- World Health Organization. (2013). *Mental health action plan 2013–2020*. http://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_R8-en.pdf
- World Health Organization. (2014). *Preventing suicide: A global imperative*. <https://apps.who.int/iris/handle/10665/131056>
- World Health Organization. (2018). *Suicide: data and figures*. <https://www.who.int/es/news-room/fact-sheets/detail/suicide>
- World Health Organization. (2021). *Live life: An implementation guide for suicide prevention in countries*. <https://www.who.int/publications/i/item/9789240026629>
- Wright, M. D. G. M. (2000). A critical-holistic paradigm for an interdependent world. *American Behavioral Scientist*, 43(5), 808–82.

How to cite this article: Sufrate-Sorzano, T., Pérez, J., Juárez-Vela, R., Garrote-Cámara, M. E., de Viñaspre, R. R., Molina-Luque, F., & Santolalla-Arnedo, I. (2022). Umbrella review of nursing interventions NIC for the treatment and prevention of suicidal behavior. *International Journal of Nursing Knowledge*, 1–12. <https://doi.org/10.1111/2047-3095.12392>